



IT'S TIME TO CLAIM VICTORY

A preventer can reduce your child's risk of an asthma attack.¹⁻³ If they're using a **blue puffer** more than twice a week, speak to your doctor.



Attack asthma and help them thrive.

Scan the QR code to visit Asthma Australia at asthma.org.au/attack

1800 ASTHMA
(1800 278 462)
asthma.org.au



References: **1.** Martinez FD, Chinchilli VM, Morgan WJ, *et al.* Use of beclomethasone dipropionate as rescue treatment for children with mild persistent asthma (TREXA): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2011; 377: 650-7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21324520>; **2.** Weiss K, Buxton M, Andersson FL *et al.* Cost-effectiveness of early intervention with once-daily budesonide in children with mild persistent asthma: results from the START study. *Pediatr Allergy Immunol* 2006; 17 Suppl 17: 21-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16573705>; **3.** Chen YZ, Busse WW, Pedersen S *et al.* Early intervention of recent onset mild persistent asthma in children aged under 11 yrs: the Steroid Treatment As Regular Therapy in early asthma (START) trial. *Pediatr Allergy Immunol* 2006; 17 Suppl 17: 7-13. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16573703>